



The mutation rate in human evolution and demographic inference

Aylwyn Scally

The germline mutation rate has long been a major source of uncertainty in human evolutionary and demographic analyses based on genetic data, but estimates have improved substantially in recent years. I discuss our current knowledge of the mutation rate in humans and the underlying biological factors affecting it, which include generation time, parental age and other developmental and reproductive timescales. There is good evidence for a slowdown in mean mutation rate during great ape evolution, but not for a more recent change within the timescale of human genetic diversity. Hence, pending evidence to the contrary, it is reasonable to use a present-day rate of approximately $0.5 \times 10^{-9} \text{ bp}^{-1} \text{ year}^{-1}$ in all human or hominin demographic analyses.

Address

Department of Genetics, University of Cambridge, Downing Street, Cambridge CB2 3EH, United Kingdom

Corresponding author: Scally, Aylwyn (aos21@cam.ac.uk)

Current Opinion in Genetics & Development 2016, 41:36-43

This review comes from a themed issue on **Genetics of human origin**

Edited by **Joshua Akey** and **Anna Di Rienzo**

<http://dx.doi.org/10.1016/j.gde.2016.07.008>

0959-437X/© 2016 Elsevier Ltd. All rights reserved.

Introduction

Population genetics provides a theoretical framework for inferring evolution, including changes in demography, based on genetic variation between individuals. It is primarily concerned with relative changes, in the sense that properties such as divergence time and population size are expressed in scaled units whose relationship to the time in years or number of individuals involved is not fixed. This is appropriate for genetic data, which are generally comparative in nature and carry no explicit record of absolute time or population size. However such data are only one of several sources of information about the evolutionary past, and the question of a timescale must be addressed if we want to relate genetic inferences to evidence from fossil, archaeological and palaeoenvironmental data.

Most demographic analyses are based on differences due to genomic mutational events, typically single-nucleotide


polymorphisms, and the quantities they estimate are naturally expressed as genetic divergence in units of substitutions per base pair. In simple terms, the genetic divergence d between two samples can be converted to a time t in years since their common ancestor by the expression $2t = d/\mu$, where μ is the mean yearly germline mutation rate over that period. Unfortunately, the question of what value of μ to use is less straightforward, as the germline mutation rate depends on multiple factors which may have varied substantially over time, and about which we may have little or no historical information. It also depends on which regions of the genome are analysed and at what level of sensitivity and specificity, making it potentially difficult to estimate an appropriate rate for a given demographic analysis or to compare estimates made using different approaches.

Mutation rates in present-day and recent human evolution

The first estimates of the human mutation rate predate the availability of molecular genetic data, and were based on the incidence of *de novo* (uninherited) disease cases where the causative allele was thought to be dominant [1,2]. In recent years, taking advantage of developments in genome sequencing technology, several new methods of estimation using genomic data have been implemented (Figure 1). Of these, estimates of the present-day genome-wide mutation rate have mostly agreed with each other, even as sequencing technologies have developed and sample sizes have grown. In particular, estimates based on whole-genome sequencing in family trios (the majority of studies) have consistently fallen in the range $1.1\text{--}1.3 \times 10^{-8} \text{ bp}^{-1}$ [3-8,9*,10,11*,12*], as did the first estimate based on identity by descent (IBD) within a pedigree [13]. Other studies have yielded slightly higher estimates however, including a more recent population-IBD estimate which obtained a value of $1.66 \times 10^{-8} \text{ bp}^{-1}$ [14], and alternative approaches using calibration against different genetic mutational processes [15,16*]. Since these methods are sensitive to somewhat older timescales than sequencing in families, which detects mutations accumulated over a generation or two at most, one possibility is that they reflect higher ancient mutation rates which have slowed in recent human evolution.

However, there are also reasons why sequencing family trios may slightly underestimate the present-day mutation rate. The main advantage of this approach is that potential samples are plentiful, allowing the measurement not just of mean rate but also variation with factors

Figure 1



Recent estimates of the human genome-wide mutation rate. Estimates are shown as yearly rates, scaled where necessary using a mean generation time of 29 years [17]; confidence intervals (90% or 95%) are shown where reported. Citation numbers and publication years are given on the x-axis. family: Family sequencing compares genomes sampled from consecutive generations in one or more families, and within each one identifies *de novo* mutations present in offspring and in neither parent [3–8,9*,10,11*,12*]. Per-generation mutation rate is calculated as the mean number of *de novo* mutations seen divided by the length of ‘callable’ genome sequenced (the number of genomic positions where a *de novo* mutation would have been called if present). IBD: Estimation based on identity by descent (IBD) detects *de novo* mutations as differences between chromosomal tracts which have been inherited IBD within or between individuals, for example in samples which are related to each other within a multi-generation pedigree. Information about the number of generations separating chromosomes may come from genealogical records [13] and/or from genetic inference [14]. aDNA: Estimation based on branch shortening in ancient DNA uses genome sequence data from an ancient human sample of known age (established with radioisotope dating) and divides the mean number of extra mutations found in present-day humans by the separation in time [18**]. PSMC: The pairwise sequential Markovian coalescent method infers ancestral effective population size from modern and ancient samples after accounting for the known age difference between them [18**]. other: Methods based on comparison with other mutational clocks: calibration using coalescent time estimates based on microsatellite mutations [15]; calibration against the recombination rate and expected variation of heterozygosity in diploid genomes [16*]. Inset: Indicative timescales over which mutations detected by each method (or which otherwise influence its estimate) have accumulated.

such as parental age and genomic distribution [8,20]. Also, unlike in other methods the temporal baseline (usually a single generation) is unambiguous. Its principal disadvantage is that single-generation *de novo* mutations are rare relative to the error rate in variant calling (60–100 mutations per individual), so false negative and false positive rates are both high and difficult to estimate. To mitigate this, genomic regions where variants are difficult to call are generally excluded via filtering, but these regions are not easy to identify and the callable genome length may be overestimated, leading to an underestimate in the per-bp mutation rate. Most of the studies cited here have attempted to quantify and account for this using simulations or validation against other methods of variant discovery, but it remains possible that true *de novo* mutation rates are consistently underestimated to some degree.

Another potential downward bias in mutation rate estimation from family trio sequencing arises from the fact that such experiments generally compare somatic cells rather than germ cells. An early post-zygotic mutation occurring before germline specification in either parent may be detected in that individual’s soma as well as in his or her offspring, and hence, seemingly present in both generations, might not be correctly identified as a *de novo* mutation [21–23]. This could be a significant factor if cellular mutation rates are particularly high in the earliest cell divisions of embryogenesis.

In principle, estimates based on IBD in a multi-generation pedigree should be less susceptible to either of these biases. Multiple accumulated mutations in IBD tracts are more easily distinguished from sequencing noise than in family sequencing, especially for larger pedigrees, and

this approach can detect all germline mutations (excepting perhaps early post-zygotic mutations in the common ancestor of a given tract). However they are not without their own methodological issues: genealogical information and uncertainty in the inference of relatedness and IBD are potential sources of error. In particular the boundaries of IBD tracts and the path of inheritance may be ambiguous, and the total extent of regions in which mutations can be detected may be quite limited except in close and inbred pedigrees. Pedigree datasets are also more difficult to collect, and since the two such genome-wide estimates published to date have not overlapped [13,14], it is difficult to assess the significance of their disagreement with other methods. We can expect forthcoming studies to help clarify this picture.

The exomic mutation rate

Mutation rates are known to vary between genomic loci [24], and the estimates discussed above are based either on whole genome data or (in the case of IBD estimates) on regions sampled genome wide without regard to location or context. Other studies, mostly using family trio sequencing, have been based on data sampled only from exomes [25–30], and have tended to yield higher values than equivalent whole-genome studies (ranging from $1.3\text{--}2.0 \times 10^{-8} \text{ bp}^{-1}$ with a mean of $1.5 \times 10^{-8} \text{ bp}^{-1}$). This is consistent with the elevated GC content of genic regions and the increased mutability of GC-rich sequence (discussed below), but it may also be that the biases discussed above are of less consequence for exome sequencing data. Consistent with the latter possibility is the fact that a recent IBD-based estimate in exomes of $1.45 \times 10^{-8} \text{ bp}^{-1}$ [31] is only slightly below the mean of trio-based estimates published so far.

Mutation rates in great ape evolution

Before the advent of high-throughput genome sequence data, estimates of the human mutation rate were generally based on phylogenetic calibration: $\mu = d/2t_s$, where d is the genetic divergence between two species and t_s the time since speciation as estimated from the fossil record. In principle, allowance must also be made for the difference between speciation and genetic divergence times, corresponding to coalescence within the ancestral population, but in practice the magnitude of this can usually only be guessed [32]. Phylogenetic calibration has some potential advantages: fossils can often be dated with relatively high accuracy using radiometric or stratigraphic methods, and since it estimates the mean substitution rate over the time separating the two species, it accounts automatically for selection and other time-varying factors which may complicate extrapolation from present-day rates.

By the time the first mutation rate estimates from *de novo* sequencing appeared, the field had largely settled on a consensus value of $1.0 \times 10^{-9} \text{ bp}^{-1} \text{ year}^{-1}$ for the yearly

rate in hominid evolution [33]. Thus the finding that *de novo* estimates were a factor of two lower than this prompted considerable debate [34]. For some events, such as the speciation of humans and chimpanzees, a higher rate had been increasingly difficult to reconcile with fossil and archaeological data, and a lower value (implying older date estimates) mostly improved concordance [35,36]. However for more ancient events a longer timescale was problematic, and to a large extent remains so still. For example, applying a present-day human mutation rate of $0.5 \times 10^{-9} \text{ bp}^{-1} \text{ year}^{-1}$ to the 2.6% genetic divergence between humans and orang-utans [37] yields an divergence time of 26 Mya. Even allowing for a large ancestral coalescent time of 5 Myr this is substantially older than the dates of 12–16 Mya typically quoted in paleoanthropological literature [38]. The difference increases for older dates: the human-macaque divergence [39] implies a speciation more than 40 Mya, whereas paleoanthropological studies generally place this node at 25–30 Mya [40].

One way to resolve this discrepancy is to regard it as evidence for a faster mutation rate 20 Mya or more, and hence a slowdown in mean rates since that time. In fact, such a hypothesis is also supported by differing branch lengths within the primates as measured from an out-group or common ancestor, with hominid (great ape) lineages being shorter than those of other primate groups by a factor of 1.4–1.6 [41*,42,43]. Because the branch shortening applies to all great apes (albeit to varying degrees), such a slowdown cannot have occurred only on the human branch, and if it occurred more recently than the hominoid ancestor (so that a higher mean mutation rate applies to dating the orang-utan divergence), it must have involved a degree of parallelism across all hominid lineages. This is not impossible for closely related species, but might be regarded as unlikely *a priori*. It is estimated that compared to humans, chimpanzees have evolved only 2% faster since divergence, and gorillas 7% faster [41*]. Indeed, a measurement of present-day mutation rate in chimpanzees based on sequencing *de novo* mutations in a multi-generation pedigree has also produced a value of $1.2 \times 10^{-8} \text{ bp}^{-1}$ [44*], very similar to equivalent human estimates.

However, timing constraints based on fossil evidence should be handled with caution. Even where fossils are themselves well dated, their correct placing relative to a particular speciation event may be far from straightforward [45,46]. There may also be important differences between the evolution of anatomical phenotypes represented in fossil taxa and the genetic differences involved in speciation, particularly when the possibility of ancestral population substructure around the time of speciation is taken into account. More fundamentally, fossil evidence tends to be more informative about lower bounds than upper bounds on speciation dates (essentially because the

presence of derived characteristics is more informative than their absence), and so 'stem' taxa which appear ancestral to a speciation event provide only weak constraints on its earliest possible date [45,47,48]. Thus it may be premature to conclude that a genetic estimate of 20–23 Mya for the orang-utan speciation is irreconcilable with fossil evidence, and the implied slowdown in mutation rate may be less than expected both in magnitude and (especially if before the orang-utan speciation) in the degree of any parallel evolution involved.

Causes and correlates of mutation rate variability

In addition to direct evidence from present-day and ancient genomic data, it may also be possible to learn about past mutation rates indirectly by studying the underlying physiological and population genetic factors affecting them. Some understanding of these factors, and how they may have varied in the past, comes from considering the cellular origins of germline mutation.

Germline mutations can arise from disruption of the DNA molecule at any time within a germline cell, but most are believed to result from errors in DNA replication during cell division, referred to as replicative mutation [49]. Over multiple generations the rate of replicative mutation will depend strongly on the mean number of cell divisions from zygote to zygote, and this can differ between species, between sexes, and perhaps also between populations due to variation in reproductive behaviour. Differences between the germline in males and females reside primarily in the sex-specific nature of gametogenesis [50], where there is a much greater number of cell divisions on the paternal lineage due to the fact that spermatogenesis involves a continuous process of stem cell division throughout adult life. This in turn contributes to a greater accumulation of *de novo* replicative mutations passed on by the father than by the mother, a phenomenon referred to as the male mutation bias and found in many species, with important evolutionary consequences [51]. Its effect in humans has been quantified in recent sequencing studies, with estimates of the male/female ratio in mean number of transmitted mutations ranging from 3.1–3.9 [8,9,12,15].

A further consequence is that the older the father, the more cell divisions his gametes will have passed through, and hence the more mutations they are likely to carry. The resulting age effect in paternally transmitted mutations has been measured at 1.2–2.0 additional *de novo* mutations per year of paternal age in recent studies [8,9,10,15,52], corresponding to a doubling from puberty to age 30. In fact this is substantially less than expected under the standard model for spermatogenesis [50,53], which predicts a factor of ten increase over the same period based on the number of cell divisions involved. Possible explanations for this discrepancy include a

revised model of spermatogenesis in which gonial stem cells pass through fewer cell divisions, or strong variation in per-cell-division mutation rates during development, with much higher rates before gametogenesis [11,23,53].

Sequencing studies initially measured no significant age effect in maternally transmitted mutations [8,9,15], consistent with replicative mutation under the longstanding reproductive model in which, after proliferation during fetal development, oocytes are held in stasis until maturation later in life and experience no postnatal cell divisions [54]. However, two more recent studies have estimated significant effects amounting to 0.51–0.86 additional mutations per year of maternal age [12,52,55]. The initial negative findings may have resulted from methodological factors (e.g. strong correlation in the data between maternal and paternal ages makes the effect difficult to discover if information on parent of origin for *de novo* mutations is not available). The distribution of parental ages sampled, which can differ even between large cohorts, particularly at the extremes of the distribution, has also been suggested as an important factor [12,53,56].

The measured maternal age effect is weaker than the paternal effect, but nevertheless supports the view that aspects of our longstanding model of gametogenesis need revision. In fact, evidence for a maternal age effect in larger-scale mutations such as chromosomal abnormalities has been available for some time [57,58], motivating the 'production line' hypothesis for oogenesis [59], which attributes the effect to a correlation between the number of pre-natal cell divisions experienced by oocytes and the age at which they are matured for ovulation. Other hypotheses have been proposed however [60], including the possibility that previously undetected post-natal oogenetic cell divisions may occur, analogous to the gametogenetic process in males [61,62]. This is supported by the discovery of germline stem cells in the ovaries of adult female humans and mice [63], and thus perhaps also by the finding of a non-zero maternal age effect in genomic mutations.

Alternatively, or in addition, non-replicative or spontaneous germline mutations may play a greater role than is generally assumed and contribute to age effects in both sexes [64]. Such mutations can arise from instability or disruption of the DNA molecule itself, for example, due to oxidative mutagens within the nucleus or exposure to ionising radiation. Unlike replicative mutation, we might expect spontaneous mutations to accumulate on germline lineages at a rate which is independent of the number of cell divisions or life history parameters such as generation time, and hence to behave more like a molecular clock. (Purely clock-like behaviour is perhaps unlikely however, as the production of oxidative mutagens is a causative factor for spontaneous mutation and is itself proportional

to metabolic rate, which also scales with generation time [65].) It has been shown that the relative contribution of spontaneous mutation depends to a large extent on the efficiency of DNA repair [49^{*}]: if such repair is rapid relative to the length of the cell cycle then most spontaneous mutations will be corrected before replication, and replicative processes will dominate. This is believed to be the case for most mutation on the human germline; however this is largely due to the correspondence between the paternal age effect and the number of mitotic divisions in males, an assumption which is perhaps undermined by the finding of a non-negligible maternal effect (assuming female post-natal mitoses are negligible).

There are also genomic loci where spontaneous mutation is expected to play a dominant role, notably CpG sites, in which the cytosine when methylated (as is usually the case in mammals [66]) is prone to spontaneous deamination from C to T. (As an aside, if spontaneous mutation contributes substantially to the maternal age effect we might expect an even stronger effect at CpG sites [49^{*}]. This was not observed in the only study so far to have examined it [12^{*},55], but larger studies in future may have greater power to detect a difference.) The more clock-like behaviour of CpG mutations is borne out in branch length comparisons within the primates and other mammals (e.g. root-to-tip distances vary by 2–4 times less than for other mutational types) [41^{*},42,67]. This makes CpG sites potentially appealing for ancestral demographic inference. However they are rare in the genome, particularly in intergenic regions (1% of sites genome-wide and 3% in exons, where they have presumably been maintained by purifying selection) [68], so their use in this way is limited to site-wise analyses ignoring haplotype information. Moreover their behaviour is not strictly clock-like but only more so than other mutations, so branch-specific factors must still be taken into account.

Discussion

The first proposed solution to the mutation rate problem was the molecular clock hypothesis of Zuckerkandl and Pauling [69], essentially a zeroth-order approximation which ignored rate variation, yet which proved surprisingly successful (in Crick's words, 'much truer than people thought at the time' [70]). In the decades since, the quest to improve upon this approximation has focused primarily on calibration against the fossil record, using increasingly sophisticated models to account for rate variation and stochasticity in fossil creation and discovery [71]. Notwithstanding the advances made in this direction, it is clear that the recent accessibility and availability of genome sequence data in humans and other species has opened a new window on the germline mutation rate.

It is also clear that generation time alone, while important, is insufficient to fully describe the dependence of mutation rates on developmental and reproductive processes.

Germline mutation depends on a plurality of related biological timescales: the ages of puberty and reproduction, the duration of fertility and of key stages in embryonic development, the cycle times of cellular processes in gametogenesis, and the efficiency of DNA repair, each potentially differing by sex or species [23,53^{**},72,73^{**}]. The sequencing studies discussed here have begun to explore these phenomena, and although some initial findings have differed or disagreed, further insights into their present-day effects and how they might have varied in the past can be expected from future sequencing on population scales. Important evidence for ancestral reproductive behaviour and life history parameters may also come from paleontological and archaeological data [74–76], and more direct evidence continues to come from ancient DNA. In particular, a recent analysis has shown that the mean generation time has not changed appreciably over at least the last 45 000 years, based on the rate of decline in linkage disequilibrium resulting from Neanderthal admixture in several ancient human samples [77^{*}].

We return therefore to the question of what mutation rate to use in analyses of human demographic evolution. Figure 1 provides a weak indication that methods sensitive to older mutation events tend to yield higher estimates, but this is somewhat confounded with potential downward bias in whole-genome estimates from family sequencing. Branch length comparisons within the apes provide no support for a substantial human-specific slowdown [41^{*}]. It may be that future developments will reveal recent modest changes in mutation rate, perhaps differing between modern human populations [78,31], driven by evolution in one or more of the factors discussed here, and possibly more substantial differences in other hominins if data become available. Pending such refinements however, a reasonable (and conservative) approach is to apply a yearly mutation rate of $0.5 \times 10^{-9} \text{ bp}^{-1} \text{ year}^{-1}$ uniformly to analyses of demographic events within or between human populations, including between modern and archaic humans.

Finally, and notwithstanding that there are many gaps in our understanding, it is worth noting that the role of the mutation rate in human demographic inference has changed markedly in recent years. Whereas genetic data were formerly regarded as definitive about topological relationships between taxa but uninformative about their time-scale, this distinction has vanished or even reversed in the case of recent human evolution. Estimates of the mutation rate have begun to converge, and it has become clear that many events in human demographic history are more complex than previously assumed, with populations diverging gradually or in convoluted ways with ongoing gene flow and admixture [79,80^{*},81]. It is fair to say that in many, perhaps even most cases, the mutation rate is no longer the principal source of ambiguity in human demographic inference.

Acknowledgements

I am grateful for support from an Isaac Newton Trust/Wellcome Trust ISSF Joint Research Grant.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Haldane J: **The rate of spontaneous mutation of a human gene.** *J Genet* 1935:226-317.
2. Nachman MW: **Haldane and the first estimates of the human mutation rate.** *J Genet* 2004, **83**:231-233.
3. Roach JC, Glusman G, Smit AFA, Huff CD, Hubley R, Shannon PT, Rowen L, Pant KP, Goodman N, Bamshad M *et al.*: **Analysis of genetic inheritance in a family quartet by whole-genome sequencing.** *Science (New York, NY)* 2010, **328**:636-639.
4. Awadalla P, Gauthier J, Myers RA, Casals F, Hamdan FF, Griffing AR, Côté M, Henrion E, Spiegelman D, Tarabeux J *et al.*: **Direct measure of the de novo mutation rate in autism and schizophrenia cohorts.** *Am J Hum Genet* 2010, **87**:316-324.
5. **1000 Genomes Project Consortium: a map of human genome variation from population-scale sequencing.** *Nature* 2010, **467**:1061-1073.
6. Conrad DF, Keebler JEM, DePristo Ma, Lindsay SJ, Zhang Y, Casals F, Idaghmour Y, Hartl CL, Torroja C, Garimella KV *et al.*: **Variation in genome-wide mutation rates within and between human families.** *Nat Genet* 2011, **43**:712-714.
7. Michaelson JJ, Shi Y, Gujral M, Zheng H, Malhotra D, Jin X, Jian M, Liu G, Greer D, Bhandari A *et al.*: **Whole-genome sequencing in autism identifies hot spots for de novo germline mutation.** *Cell* 2012, **151**:1431-1442.
8. Kong A, Frigge ML, Masson G, Besenbacher S, Sulem P, Magnusson G, Gudjonsson Sa, Sigurdsson A, Jonasdottir A, Jonasdottir A *et al.*: **Rate of de novo mutations and the importance of father's age to disease risk.** *Nature* 2012, **488**:471-475.
9. Consortium of the Netherlands Consortium: **Whole-genome sequence variation, population structure and demographic history of the Dutch population.** *Nat Genet* 2014, **46**:818-825.
- The first sequencing study of de novo mutations in a cohort comprising hundreds of parent-offspring trios.
10. Besenbacher S, Liu S, Izarzugaza JMG, Grove J, Belling K, Bork-Jensen J, Huang S, Als TD, Li S, Yadav R *et al.*: **Novel variation and de novo mutation rates in population-wide de novo assembled Danish trios.** *Nat Commun* 2015, **6**:5969.
11. Rahbari R, Wuster A, Lindsay SJ, Hardwick RJ, Alexandrov LB, Al Turki S, Dominiczak A, Morris A, Porteous D, Smith B *et al.*: **Timing, rates and spectra of human germline mutation.** *Nat Genet* 2015: 1-11.
- Presents the results of whole genome de novo mutation sequencing of multi-sibling families, enabling the authors to analyse shared mutations and estimate rates of germline mosaicism and mutation recurrence.
12. Wong WSW, Solomon BD, Bodian DL, Kothiyal P, Eley G, Huddleston KC, Baker R, Thach DC, Iyer RK, Vockley JG *et al.*: **New observations on maternal age effect on germline de novo mutations.** *Nat Commun* 2016, **7**:10486.
- The largest whole-genome sequencing study of de novo mutations to date, revealing a maternal age effect in transmitted mutations.
13. Campbell CD, Chong JX, Malig M, Ko A, Dumont BL, Han L, Vives L, O'Roak BJ, Sudmant PH, Shendure J *et al.*: **Estimating the human mutation rate using autozygosity in a founder population.** *Nat Genet* 2012, **44**:1277-1281.
14. Palamara PF, Francioli LC, Wilton PR, Genovese G, Gusev A, Finucane HK, Sankararaman S, Sunyaev SR, De Bakker PIW, Wakeley J *et al.*: **Leveraging distant relatedness to quantify human mutation and gene-conversion rates.** *Am J Hum Genet* 2015, **97**:775-789.
15. Sun JX, Helgason A, Masson G, Ebenesersdóttir SS, Li H, Mallick S, Gnerre S, Patterson N, Kong A, Reich D *et al.*: **A direct characterization of human mutation based on microsatellites.** *Nat Genet* 2012, **44**:1161-1165.
16. Lipson M, Loh PR, Sankararaman S, Patterson N, Berger B, Reich D: **Calibrating the human mutation rate via ancestral recombination density in diploid genomes.** *PLoS Genet* 2015, **11**:e1005550.
- Presents an interesting method of mutation rate estimation based on calibration against the (better known) recombination clock and simulations of the distribution of genomic polymorphisms as a function of genetic distance.
17. Fenner JN: **Cross-cultural estimation of the human generation interval for use in genetics-based population divergence studies.** *Am J Phys Anthropol* 2005, **128**:415-423.
18. Fu Q, Li H, Moorjani P, Jay F, Slepchenko SM, Bondarev Aa, Johnson PLF, Aximu-Petri A, Prüfer K, de Filippo C *et al.*: **Genome sequence of a 45,000-year-old modern human from western Siberia.** *Nature* 2014, **514**:445-449.
- An important ancient human sample studied using ancient DNA, presenting two novel approaches to estimating the mutation rate in recent human evolution.
19. Li H, Durbin R: **Inference of human population history from individual whole-genome sequences.** *Nature* 2011, **475**:493-496.
20. Francioli LC, Polak PP, Koren A, Menelaou A, Chun S, Renkens I, van Duijn CM, Swertz M, Wijmenga C, van Ommen G *et al.*: **Genome-wide patterns and properties of de novo mutations in humans.** *Nat Genet* 2015, **47**:822-826.
21. Campbell IM, Yuan B, Robberecht C, Pfundt R, Szafranski P, McEntagart ME, Nagamani SC, Erez A, Bartnik M, Wiśniowiecka-Kowalnik B *et al.*: **Parental somatic mosaicism is underrecognized and influences recurrence risk of genomic disorders.** *Am J Hum Genet* 2014, **95**:173-182.
22. Acuna-Hidalgo R, Bo T, Kwint MP, van de Vorst M, Pinelli M, Veltman JA, Hoischen A, Vissers LE, Gilissen C: **Post-zygotic point mutations are an underrecognized source of de novo genomic variation.** *Am J Hum Genet* 2015, **97**:67-74.
23. Scally A: **Mutation rates and the evolution of germline structure.** *Phil Trans R Soc B: Biol Sci.* 2016:371.
24. Tyekucheva S, Makova KD, Karro JE, Hardison RC, Miller W, Chiaromonte F: **Human-macaque comparisons illuminate variation in neutral substitution rates.** *Genome Biol* 2008, **9**:R76.
25. Sanders SJ, Murtha MT, Gupta AR, Murdoch JD, Raubeson MJ, Willsey AJ, Ercan-Sencicek aG, DiLullo NM, Parikshak NN, Stein JL *et al.*: **De novo mutations revealed by whole-exome sequencing are strongly associated with autism.** *Nature* 2012, **485**:237-241.
26. O'Roak BJ, Vives L, Girirajan S, Karakoc E, Krumm N, Coe BP, Levy R, Ko A, Lee C, Smith JD *et al.*: **Sporadic autism exomes reveal a highly interconnected protein network of de novo mutations.** *Nature* 2012, **485**:246-250.
27. Iossifov I, Ronemus M, Levy D, Wang Z, Hakker I, Rosenbaum J, Yamrom B, Lee Yh, Narzisi G, Leotta A *et al.*: **De novo gene disruptions in children on the autistic spectrum.** *Neuron* 2012, **74**:285-299.
28. Neale BM, Kou Y, Liu L, Ma'ayan A, Samocha KE, Sabo A, Lin CF, Stevens C, Wang LS, Makarov V *et al.*: **Patterns and rates of exonic de novo mutations in autism spectrum disorders.** *Nature* 2012, **485**:242-245.
29. Zaidi S, Choi M, Wakimoto H, Ma L, Jiang J, Overton JD, Romano-Adesman A, Bjornson RD, Breitbart RE, Brown KK *et al.*: **De novo mutations in histone-modifying genes in congenital heart disease.** *Nature* 2013, **498**:220-223.
30. Fromer M, Pocklington AJ, Kavanagh DH, Williams HJ, Dwyer S, Gormley P, Georgieva L, Rees E, Palta P, Ruderfer DM *et al.*: **De novo mutations in schizophrenia implicate synaptic networks.** *Nature* 2014, **506**:179-184.
31. Narasimhan VM, Rahbari R, Scally A, Wuster A, Mason D, Xue Y, Wright J, Trembath RC, Maher ER, van Heel DA *et al.*: **A direct multi-generational estimate of the human mutation rate from**

- autozygous segments seen in thousands of parentally related individuals.* bioRxiv 2016.
32. Burgess R, Yang Z: *Estimation of hominoid ancestral population sizes under Bayesian coalescent models incorporating mutation rate variation and sequencing errors.* Mol Biol Evol 2008, **25**:1979-1994.
 33. Nachman MW, Crowell SL: *Estimate of the mutation rate per nucleotide in humans.* Genetics 2000, **156**:297-304.
 34. Scally A, Durbin R: *Revising the human mutation rate: implications for understanding human evolution.* Nat Rev Genet 2012, **13**:745-753.
 35. Scally A, Dutheil JY, Hillier LW, Jordan GE, Goodhead I, Herrero J, Hobolth A, Lappalainen T, Mailund T, Marques-Bonet T *et al.*: *Insights into hominid evolution from the gorilla genome sequence.* Nature 2012, **483**:169-175.
 36. Hawks J: **Longer time scale for human evolution.** *Proc Natl Acad Sci U S A.* 2012:1-2.
 37. Locke DP, Hillier LW, Warren WC, Worley KC, Nazareth LV, Muzny DM, Yang SP, Wang Z, Chinwalla AT, Minx P *et al.*: *Comparative and demographic analysis of orang-utan genomes.* Nature 2011, **469**:529-533.
 38. Hartwig W (Ed): *The Primate Fossil Record.* Cambridge University Press; 2002.
 39. Carbone L, Alan Harris R, Gnerre S, Veeramah KR, Lorente-Galdos B, Huddleston J, Meyer TJ, Herrero J, Roos C, Aken B *et al.*: *Gibbon genome and the fast karyotype evolution of small apes.* Nature 2014, **513**:195-201.
 40. Stevens NJ, Seiffert ER, O'Connor PM, Roberts EM, Schmitz MD, Krause C, Gorscak E, Ngasala S, Hieronymus TL, Temu J: *Palaeontological evidence for an Oligocene divergence between Old World monkeys and apes.* Nature 2013, **497**:611-614.
 41. Moorjani P, Amorim CEG, Arndt PF, Przeworski M: *Variation in the molecular clock of primates.* bioRxiv 2016:036434.
An up-to-date phylogenetic analysis of primate whole genome alignments, quantifying the extent of branch length variation and its relationship to life history and other factors.
 42. Kim SH, Elango N, Warden C, Vigoda E, Yi SV: *Heterogeneous genomic molecular clocks in primates.* PLoS Genet 2006, **2**:e163.
 43. Steiper ME, Young NM, Sukarna TY: *Genomic data support the hominoid slowdown and an Early Oligocene estimate for the hominoid-cercopithecoid divergence.* Proc Natl Acad Sci U S A 2004, **101**:17021-17026.
 44. Venn O, Turner I, Mathieson I, de Groot N, Bontrop R, McVean G: *Strong male bias drives germline mutation in chimpanzees.* Science 2014, **344**:1272-1275.
The first measurement of mutation rate (and age effects) in a non-human primate, also provides a good example of the methodology involved in sequencing *de novo* mutations in a small pedigree.
 45. Steiper ME, Young NM: *Timing primate evolution: lessons from the discordance between molecular and paleontological estimates.* Evol Anthropol: Iss News Rev 2008, **17**:179-188.
 46. Jensen-Seaman MI, Hooper-Boyd KA: *Molecular clocks: determining the age of the human-chimpanzee divergence.* eLS 2013.
 47. Ho SY, Phillips MJ: *Accounting for calibration uncertainty in phylogenetic estimation of evolutionary divergence times.* Syst Biol 2009:syp035.
 48. Pozzi L, Hodgson JA, Burrell AS, Disotell TR: *The stem catarrhine saadanius does not inform the timing of the origin of crown catarrhines.* J Hum Evol 2011, **61**:209-210.
 49. Gao Z, Wyman MJ, Sella G, Przeworski M: *Interpreting the dependence of mutation rates on age and time.* PLOS Biol 2016, **14**:e1002355.
An insightful analysis of replicative and non-replicative modes of germline mutation and the impact of DNA repair and other timescales on their relative contribution.
 50. Vogel F, Motulsky A: *Human Genetics: Problems and Approaches.* Berlin: Springer; 1979.
 51. Ellegren H: *Characteristics, causes and evolutionary consequences of male-biased mutation.* Proc R. Soc B: Biol Sci 2007, **274**:1-10.
 52. McRae JF, Clayton S, Fitzgerald TW, Kaplanis J, Prigmore E, Rajan D, Sifrim A, Aitken S, Akawi N, Alvi M *et al.*: *Prevalence, phenotype and architecture of developmental disorders caused by *de novo* mutation.* bioRxiv 2016:049056.
 53. Ségurel L, Wyman MJ, Przeworski M: *Determinants of mutation rate variation in the human germline.* Annu Rev Genom Hum Genet 2014, **15**:47-70.
An excellent comprehensive review of genomic mutation in humans and the underlying factors affecting it.
 54. Zuckerman S: *The number of oocytes in the mature ovary.* Recent Prog Horm Res 1951, **6**:63-109.
 55. Goldmann JM, Wong WS, Pinelli M, Farrah T, Bodian D, Stittrich AB, Glusman G, Vissers LE, Hoischen A, Roach JC *et al.*: *Parent-of-origin-specific signatures of *de novo* mutations.* Nat Genet 2016.
 56. Forster P, Hohoff C, Dunkelmann B, Schurenkamp M, Pfeiffer H, Neuhuber F, Brinkmann B: *Elevated germline mutation rate in teenage fathers.* Proc R Soc B: Biol Sci 2015, **282**:20142898.
 57. Ferguson-Smith M, Yates J: *Maternal age specific rates for chromosome aberrations and factors influencing them: report of a collaborative European study on 52 965 amniocenteses.* Prenatal Diagn 1984, **4**:5-44.
 58. Pellestor F, Andréo B, Arnal F, Humeau C, Demaille J: *Maternal aging and chromosomal abnormalities: new data drawn from *in vitro* unfertilized human oocytes.* Hum Genet 2003, **112**:195-203.
 59. Henderson S, Edwards R: *Chiasma frequency and maternal age in mammals.* Nature 1968.
 60. Perry JR, Murray A, Day FR, Ong KK: *Molecular insights into the aetiology of female reproductive ageing.* Nat Rev Endocrinol 2015.
 61. Woods DC, Telfer EE, Tilly JL: *Oocyte family trees: old branches or new stems?* PLoS Genet 2012, **8**:e1002848.
 62. Dunlop CE, Telfer EE, Anderson Ra: *Ovarian germline stem cells.* Stem Cell Res Ther 2014, **5**:98.
 63. White YA, Woods DC, Takai Y, Ishihara O, Seki H, Tilly JL: *Oocyte formation by mitotically active germ cells purified from ovaries of reproductive-age women.* Nat Med 2012, **18**:413-421.
 64. Stoltzfus A: *Evidence for a predominant role of oxidative damage in germline mutation in mammals.* Mutat Res/Fundam Mol Mech Mutagen 2008, **644**:71-73.
 65. Gillooly JF, McCoy MW, Allen AP: *Effects of metabolic rate on protein evolution.* Biol Lett 2007, **3**:655-659.
 66. Bird AP: *DNA methylation and the frequency of CPG in animal DNA.* Nucleic Acids Res 1980, **8**:1499-1504.
 67. Hwang DG, Green P: **Bayesian Markov chain Monte Carlo sequence analysis reveals varying neutral substitution patterns in mammalian evolution.** *Proc Natl Acad Sci U S A.* . 2004:13994-14001.
 68. Saxonov S, Berg P, Brutlag DL: *A genome-wide analysis of CPG dinucleotides in the human genome distinguishes two distinct classes of promoters.* Proc Natl Acad Sci U S A 2006, **103**:1412-1417.
 69. Zuckerkandl E, Pauling L: *Evolutionary divergence and convergence in proteins.* Evol Genes Proteins 1965, **97**:97-166.
 70. Crick F: **The impact of linus pauling on molecular biology.** Edited by Krishnamurty RS. The Pauling Symposium: A Discourse on the Art of Biography . Corvallis, OR: Oregon State University Libraries; 1996.
 71. dos Reis M, Donoghue PC, Yang Z: *Bayesian molecular clock dating of species divergences in the genomics era.* Nat Rev Genet 2016, **17**:71-80.
 72. Thomas GWC, Hahn MW: *The human mutation rate is increasing, even as it slows.* Mol Biol Evol 2014, **31**:253-257.

73. Amster G, Sella G: **Life history effects on the molecular clock of autosomes and sex chromosomes**. *Proc Natl Acad Sci U S A* . 2016:1588-1593.
- Analysis of the issues surrounding life history effects on mutation time, focusing in particular on sex differences, considering the implications for genomic variation between chromosomes and for phylogenetic analyses of speciation.
74. Robson SL, Wood B: *Hominin life history: reconstruction and evolution*. *J Anat* 2008, **212**:394-425.
75. Smith TM, Tafforeau P, Reid DJ, Pouech J, Lazzari V, Zermeno JP, Guatelli-Steinberg D, Olejniczak AJ, Hoffman A, Radovic J *et al.*: **Dental evidence for ontogenetic differences between modern humans and Neanderthals**. *Proc Natl Acad Sci U S A* . 2010:20923-20928.
76. Gluckman PD, Hanson MA: *Evolution, development and timing of puberty*. *Trends Endocrinol Metab* 2006, **17**:7-12.
77. Moorjani P, Sankararaman S, Fu Q, Przeworski M, Patterson N, Reich D: **A genetic method for dating ancient genomes provides a direct estimate of human generation interval in the last 45,000 years**. *Proc Natl Acad Sci U S A* . 2016:201514696.
- Calibrates the decay of archaic-admixture-associated linkage disequilibrium in several ancient human genomes against their known ages, providing direct evidence for constancy of the generation time in recent human evolution.
78. Harris K: *Evidence for recent, population-specific evolution of the human mutation rate*. *Proc Natl Acad Sci U S A* 2015, **112**:3439-3444.
79. Reich D, Patterson N, Kircher M, Delfin F, Nandineni MRR, Pugach I, Ko AMSMS, Ko YC, Jinam TAA, Phipps MEE *et al.*: *Denisova admixture and the first modern human dispersals into southeast Asia and Oceania*. *Am J Hum Genet* 2011, **89**:516-528.
80. Pickrell JK, Reich D: *Toward a new history and geography of human genes informed by ancient DNA*. *Trends Genet* 2014, **30**:377-389.
- A good survey of recent developments in human evolutionary genomics and discussion of the power and potential of ancient DNA. Reviews the evidence for complex events and relationships between populations in human demographic history.
81. Schiffels S, Durbin R: *Inferring human population size and separation history from multiple genome sequences*. *Nat Genet* 2014, **46**:919-925.